BD Vacutainer® System BD Diagnostics - Preanalytical Systems



Tube Guide including Order of Draw*

Please display this in your clinical areas beside your venepuncture equipment

Blood samples should be taken in the following order:				
Cat. No. & Draw Vol.	Colour Code	Tube Type	Determinations	
		Blood Culture	Blood Cultures	
Draw Volume	Light Blue	Sodium Citrate	Coagulation Tests	
		ECD		
Draw Volume	Black			
Draw Volume	Red	Serum	eg Theraputic Drug Monitoring	
Draw Volume	Orange	1131		
Draw Volume	Gold	SST™II	eg Lytes, LFTs, Iron, Hormones	
Draw Volume	Green	Heparin & PST™II	eg. Lactate, Ammonia, Plasma K	
Draw Volume	Lavender	EDTA	eg. CBC, Retics, ESR, CD4	
Draw Volume	Pearlescent	PPT	Use as discard	
Draw Volume	Pink	Cross Match	Tranfusion Medicine	
Draw Volume	Grey	Fluoride Oxalate	eg. Glucose, glucose Tolerence	
Draw Volume	Royal Blue	Trace Element	eg. Lead, Zinc, Chromium, cobalt	

BD Diagnostics - Preanalytical Systems
The Danby Building, Edmund Halley Road, Oxford Science Park, Oxford, OX4 4DQ Tel. No.: 01865 781603 Fax: 01865 781528

BD Vacutainer® System BD Diagnostics - Preanalytical Systems



Mixing Guidelines

All BD Vacutainer tubes require immediate mixing following collection

Colour Code	Tube Type	Inversions
Light Blue	Sodium Citrate	3-4 Times
	Carlings Cityata FCD	0 10 Times
Black	Joanam entrate 25%	o ro riines
	DCT™	F. C. Timore
Orange		5 6 1111105
Red	Serum	5-6 Times
Gold	SST™ II	5-6 Times
Green	Heparin & PST™ II	8-10 Times
Lavender	EDTA	8-10 Times
Pearlescent	РРТ	8-10 Times
Pink	Cross Match	8-10 Times
Grey	Fluoride Oxalate	8-10 Times
Royal Blue	Trace Element	8-10 Times

Insufficient mixing can result in inaccurate test results and the need to re-draw